

DECSO emission versions

Authors: Ronald van der A, Jieying Ding, Bas Mijling, Xiaojuan Lin, October 2025

The DECSO algorithm (Daily Emission estimation Constrained by Satellite Observations) has been applied successfully for NO_x emission estimation in East Asia, Middle East, South Africa, and India. As the algorithm is further improved, emission estimations are updated accordingly. Below you can find an overview of the different algorithm versions. An extensive algorithm description can be found in Mijling and Van der A (2012), and the GlobEmission ATBD.

Version 1

- CHIMERE V2006
- No biogenic emissions
- INTEX-B emission inventory
- Landuse by GLCF database (1993)
- European diurnal cycle
- Boundary conditions LMDzINCA (gas), GOCART (aerosol)

Used in:

- Mijling and Van der A (2012)
- GlobEmission, phase 1 (East Asia)

Version 2

- Modelling CHIMERE with sector-dependent emission injection heights
- Fast back-trajectory calculations
- Adjusted retrieval error tropospheric NO₂
- Update of NO_x-correlated pollutants
- Noise and bias reduction in remote areas
- Full Kalman matrix inversion using LDL decomposition
- Initial emission inventories:
 - South Africa: EDGAR v4.2, 2008
 - India: EDGAR v4.2, 2008

Used in:

- Mijling et al. (2013)
- GlobEmission, phase 1 (India, South Africa)

Version v3a

- CHIMERE V2006.
- Initial emission inventories:
 - East Asia: MEIC 2008 (China) + INTEX-B (outside China)
 - South Africa: EDGAR v4.2, 2008
 - Middle East: EDGAR v4.2, 2008
- Diurnal cycle: flattened for China, European for other regions.
- Calculation speed: switching from g95 to ifort compiler, and calculating matrix inversions with LAPACK

Used in:

- GlobEmission, phase 2 (East Asia, Middle East, South Africa)

The emission estimates show unrealistic day-to-day (and possibly month-to-month) fluctuations of emissions. Emission noise in low-emitting areas (introduction of positive emission bias, and unrealistic seasonal cycle when assimilating OMI measurements).

Version v3b

- CHIMERE V2013: new transport schemes, secondary organic aerosol chemistry, updated chemical reaction rates.
- New land use data: GlobCover Land cover (2009).
- Biogenic emissions by MEGAN
- Initial emission inventories:
 - East Asia: MEIC 2010 (China) + INTEX-B (outside China), regridded.
 - Middle East: HTAP v2 (EDGAR v4.3)
- Reduction of day-to-day emission fluctuations by OmF criterium [-5,10].
- Diurnal cycle: flattened for Middle East, European for East Asia.

Used in:

- Ding et al. (2015)
- GlobEmission, phase 2 (East Asia, Middle East)

Version 4

- Reduction of day-to-day emission fluctuations by 3-sigma (emission error) criterion
- New parametrization of R matrix
- Diurnal cycle: flattened
- OmF criterion has been removed

Used in:

- GlobEmission, phase 2 (East Asia)

The emission noise is greatly reduced. The regional emission totals go down, but this is mainly related to the emission noise reduction (reduction of positive bias). Individual hot-spots, especially when undersampled, can disappear (e.g. Ulaanbaatar, and some power plants in North-East China). A slower convergence rate at changing emission signals. But 10% more observations have been used in this version compared with DECSO v3b.

Version 5

- Switch off biogenic emissions from MEGAN in CHIMERE, estimate total surface emissions instead of anthropogenic emissions
- Change the threshold of the sensitivity matrix from 0.05 h to 0.1 h.

Used in:

- GlobEmission, phase 2 (East Asia)
- Ding et al. (2017a, 2017b)

The total emissions are more realistic and we reduce the uncertainties in biogenic emissions from CHIMERE. The change of H threshold helps to reduce biases coming from the uncertainties occurring at the edge of a NO₂ plume.

This version is extensively validated for East China in Ding et al. (2017b)

Version 5.1

- Set maritime inject height of newly-found maritime at 40 m.
- Exclude the observations with a large pixel size by filtering out 8 pixels at each side of the swath
- Exclude the observations with a cloud fraction larger than 50%

Used in:

- Ding et al. (2018)
- For QA4ECV data: Ding et al. (2022)

With the new settings, we get more clear shipping tracks near Chinese coast areas.

Version 5.2-TROPOMI

- Results are generated on 0.125 degree resolution for a smaller domain.
- Timestep of the CHIMERE decreased to 7.5 min.
- The lifetime fit of NO_x has been optimized for more precise local lifetimes.
- New parametrization of the R-matrix for TROPOMI
- The value of the minimum of H-elements has been set to 0.05 to avoid amplification of noise in the inverse calculation.
- The correlation length of Q is set to 1 km.
- Improved regularisation of the inverse calculation of (KSK+R).
- The NO₂ climatology for the free troposphere has been corrected.

Used in

- Van der A et al. (2020)

This version is developed for using TROPOMI NO₂ data. Since TROPOMI observations have a much higher resolution, we can also switch to higher resolution output.

Version 5.2-TROPOMI-superobservations

- In this variant of version 5.2 (1) TROPOMI data have been regridded into super-observations before using as input and (2) Results are generated on 0.25 degree resolution.
- Because of the different resolution, different parametrizations are used for the R-matrix and Q_matrix that describe the errors due to the model, measurements and representation.

Used in:

- Ding et al. (2020)

Since this version use super observations and generates output on the lower resolution of 0.25 degree it is much faster and can easily calculate emissions for large regions.

Version 5.5-TROPOMI-superobservations

- In this version superobservations are generated for columns till 500 hPa to remove upper-tropospheric effects.
- No error correction for negative obs.
- Correlation length of emissions is 2 km (i.o. 10 km)
- Radius of taking correlation length into account is 50 km (i.o. 300 km)
- Shorter steps in lifetime fit to reach more often convergence (dk=0.2k instead of 0.5k.)
- Lifetime fit: if the last iteration is resulting in a worse cost function, then take the previous iteration.
- New landuse data base: Copernicus Landcover 2019.
- Use of PAL data for TROPOMI NO₂ data

Used in:

- Used as SEEDS deliverable for Europe

Version 6.0-TROPOMI

- superobservations till 500 hPa are used
- CHIMERE 2020r3 is replacing th older CHIMERE version
- Time resolution step (timeres=8) of 7.5 minutes

Used in:

- Used as SEEDS deliverable for High-Resolution output (Lowlands)

Version 6.1-TROPOMI (not released)

- Combined NH₃ and NO_x inversion. (see Ding et al., 2024)
- maximum distance in H is 200 km.
- new start -field HTAP3-2018
- Minimum value of H (H_min) becomes 0.01 (was 0.05)
- 3-sigma limit turned off

Used in:

- Used in X. Zhang et al. (2023)

Version 6.2-TROPOMI

- Timeres=8 for 0.2x0.2 ° (default), timeres=12 for 0.1x0.15 ° (Spain) and timeres=30 for 0.05 x0.05 (lowlands)
- The range *Hdistance_max* of the sensitivity function H is reduced to 150 km for 0.2x0.2° (default), and 50-100 km for lower resolutions (Spain, Lowlands)
- Number of iterations for the lifetime fit is limited to reduce the calculation time.
- R is based on the level2 error estimates (about.0.3), $Q_{abs}=0.025$, $Q_{rel}=0.02$
- The subgridding over a satellite pixel is based on 16x16 subgrids (instead of 32x8)
- The NO₂ observations are based on TROPOMI retrieval version 2.4. For the highest resolution of 0.05° no superobservations are used.
- In the calculated emissions negative emission are set to zero without compensation the neighbouring grid cells.
- Soil emission of NO_x are provided based on a fitted seasonal parametrisation.
- TROPOMI version 2.4 has been used

Used in:

- Used as SEEDS deliverable for Europe

Version 6.2HR (not released)

Same as version 6.2 except for the HR grid and:

- Applied to normal TROPOMI observations (i.e. no superobservations used)
- The resolution of the grid is 0.05x0.05 degree
- A reduced number of observations (less than 4000) is used per day
- More model steps in trajectories. N=30
- Because of calculation speed: no iterations for the lifetime fit

Version 6.3 and 6.3.1

Inversion for NO_x and NH₃ emissions:

- New boundary conditions in CHIMERE: LMDzINCA_2010_2017_V2020
- New superobservations including border values (accidentally empty in previous versions)
- Version 6.3.1 includes a new split up in soil NO_x emissions and anthropogenic NO_x emissions (fitted per month and land-use type instead of a seasonal parametrization)

Version 6.3.1 is published in van der A et al., 2024, Ding et al., 2024, and Lin et al., 2024.

Version 6.4 and 6.4.1

- New vertical emission profiles (based on Bieser et al., 2013)
- Corrected upper-tropospheric/stratospheric correction of the TROPOMI NO₂ observations (via CTM-ANA priori files)
- 6.4.1 includes the soil-anthropogenic split-up

Version 6.5

- Better integration of parallel processing of the NO_x and NH₃ emissions
- Option for reference sector correction for biases in the NO₂ observations.
- Correction of bug in super-observations to get the correct ceiling pressure level.
- Improved accuracy calculation for the TROPOMI NO₂ super-observations.
- User-friendly options for different meteo domain and modelling domain
- Seasonal cycle for VOC emissions included
- Emission height profile for ship emissions adapted
- Error correction for the high concentrations (so the assimilation favors high concentrations)

Version 6.5 is used in ESA projects DECSO-NRT and AGATE and EU-Horizon project GreenEO.

Version 6.5-HR (only for the Netherlands and region around Shanghai)

- Basic correction for the level2 NO₂ observations in wintertime for winds from Northerly to Westerly directions for the Netherlands
- A fixed lifetime (3 hours) is used in the sensitivity calculation.

References

Lin et al., European soil NO_x emissions derived from satellite NO₂ observations, *Journal of Geophysical Research – Atmospheres*, submitted, 2024

Ding, J., van der A, R., Eskes, H., Damers, E., Shephard, M., Wichink Kruit, R., Guevara, M., and Tarrason, L.: Ammonia emission estimates using CrIS satellite observations over Europe, *Atmos. Chem. Phys.*, 24, 10583–10599, <https://doi.org/10.5194/acp-24-10583-2024>, 2024.

van der A, R. J., Ding, J., and Eskes, H.: Monitoring European anthropogenic NO_x emissions from space, *Atmos. Chem. Phys.*, 24, 7523–7534, <https://doi.org/10.5194/acp-24-7523-2024>, 2024.

Zhang, X., van der A, R., Ding, J., Zhang, X., and Yin, Y.: Significant contribution of inland ships to the total NO_x emissions along the Yangtze River, *Atmos. Chem. Phys.*, 23, 5587–5604, <https://doi.org/10.5194/acp-23-5587-2023>, 2023.

Jieying Ding, Ronald van der A, Bas Mijling, Jos de Laat, Henk Eskes, K. Folkert Boersma, NO_x emissions in India derived from OMI satellite observations, *Atmospheric Environment: X*, 14, 100174, <https://doi.org/10.1016/j.aeoa.2022.100174>, 2022.

Ding, J., van der A, R. J., Eskes, H. J., Mijling, B., Stavrou, T., van Geffen, J. H. G. M., Veeffkind, J.P., NO_x emissions reduction and rebound in China due to the COVID-19 crisis, *Geophysical Research Letters*, 46, e2020GL089912, <https://doi.org/10.1029/2020GL089912>, 2020

van der A, R.J., de Laat, A.T.J., Ding, J., Eskes, H.J., Connecting the dots: NO_x emissions along a West Siberian natural gas pipeline, *npj. Clim. Atmos. Sci.* 3, 16, <https://doi.org/10.1038/s41612-020-0119-z>, 2020

Ding, J. van der A, R. J., Mijling, B., Jalkanen, J.-P. Johansson, L., and Levelt, P. F.: Maritime NO_x emissions over Chinese seas derived from satellite observations, *Geophysical Research Letters*, 45, <https://doi.org/10.1002/2017GL076788>, 2018

Ding, J., van der A, R. J., Mijling, B., and Levelt, P. F.: Space-based NO_x emission estimates over remote regions improved in DECSO, *Atmos. Meas. Tech.*, 10, 925-938, <https://doi.org/10.5194/amt-10-925-2017>, 2017a.

Ding, J., Miyazaki, K., van der A, R.J., Mijling, B., Kurokawa, J., Cho, S., Janssens-Maenhout, G., Zhang, Q., Liu, F., and Levelt, P.F., Intercomparison of NO_x emission inventories over East Asia, *Atm. Chem. Phys.*, 17, 10125-10141, [doi:doi.org/10.5194/acp-17-10125-2017](https://doi.org/10.5194/acp-17-10125-2017), 2017b.

Ding, J., van der A, R. J., Mijling, B., Levelt, P. F., and Hao, N.: NO_x emission estimates during the 2014 Youth Olympic Games in Nanjing, *Atmos. Chem. Phys.*, 15, 9399-9412, [doi:10.5194/acp-15-9399-2015](https://doi.org/10.5194/acp-15-9399-2015), 2015.

Mijling, B. and R. van der A, Using daily satellite observations to estimate emissions of short-lived air pollutants on a mesoscopic scale. *J. Geophys. Res.* 117, D17, [doi:10.1029/2012JD017817](https://doi.org/10.1029/2012JD017817), 2012.

Mijling, B., R.J. van der A, R. J., and Q. Zhang, Regional nitrogen oxides emission trends in East Asia observed from space, *Atmos. Chem. Phys.* 13, 12003-12012, [doi:10.5194/acp-13-12003-2013](https://doi.org/10.5194/acp-13-12003-2013), 2013.

SOFTWARE and KEY DATA of Version 6.1

Software/data	Version	Ref
Inverse algorithm	DECSO v6.4	
Chemical Transport Model	CHIMERE v2020r3	
NO2 observations	TROPOMI v.2.4	
Meteorological data	ECMWF operational model	
Land-use database	Copernicus LandCover	
Start field emissions	HTAP v3	

PARAMETERS in Version 6.2

Parameter	Value
Time resolution transport kernel	7.5 minutes or less
Initial lifetime	6 hours
Initial maximum lifetime	6 hours
Maximum lifetime during iterations	100 hours
Minimum Lifetime	0.5 hour
Maximum Lifetime	16 hours
Smallest allowed value for matrix H elements	0.01
Alpha (Regularisation factor for lifetime fit)	0.01
Emission correlation length	0.5 km
Q _{abs} Absolute error of emissions	Depends on resolution
Q _{rel} Relative error of emissions	Depends on resolution
OmF correlation length (default)	10 km
Pa _{radius} (Max. distance for cov. analysis)	50 km
absMsigma (absolute error of σ_R)	0.1
relMsigma (relative error of σ_R)	0.05
Tikhonov _{threshold}	0.01
Error of start-field Pa of emissions on day 1	0.3 [1e15 molec/cm ² /h]
Maximum distance for emission-concentration relationship	150 km or less for higher resolution emissions

Version	Year of release	Study using this data version
1	2012	Mijling and Van der A (2012)
2	2013	Mijling et al. (2013)
3a	2014	GlobEmission, phase 2 (East Asia, Middle East, South Africa)
3b	2015	Ding et al. (2015), GlobEmission, phase 2 (East Asia, Middle East)
4	2016	GlobEmission, phase 2 (East Asia)
5.0	2017	Ding et al. (2017a, 2017b)
5.1	2018	Ding et al. (2018)
5.2	2019	Van der A et al. (2020)
5.2-superobs	2020	Ding et al. (2020)
5.6	2022	SEEDS, first phase, 0.2x0.2 degree
6.0	2022	SEEDS, first phase, 0.05x0.05
6.1	2022	Zhang et al. (2023)
6.2	2023	SEEDS, second phase
6.3	2024	van der A et al., 2024, Ding et al., 2024, and Lin et al., 2024.
6.4	2024	CAMEO project